



第22回トポロジカル物質科学セミナー Topological Material Science Seminar (22)

Anomalous Hall effects in Chiral Superfluids

Prof. James A. Sauls

Department of Physics & Astronomy
Northwestern University, Evanston, USA

Place: Room 525, Department of Physics, Graduate School of Science,
Building No.5, Kyoto University

Date : November 4 (Friday), 2016

Time: 10:30-

Abstract:

The superfluid phases of ^3He are paradigms for spontaneous symmetry breaking in quantum field theory and condensed matter physics. The microscopic physics underlying the phenomenology of ^3He - that of an interacting Fermi system with strong-coupling between Fermions to *paramagnons* - provided the basic theoretical model for spin-triplet superconductivity in Sr_2RuO_4 [1]. The phases of bulk superfluid ^3He are also paradigms for topological order and the subject of intense theoretical and experimental research [2]. I discuss signatures of broken space-time symmetries - particularly, parity and time-reversal (BTRP) - and the implications for topological order of chiral superfluids. I highlight signatures of BTRP in $^3\text{He-A}$ [3], and chiral superconductors [4]. I summarize the theory for the anomalous Hall effect for electron transport in chiral superfluids, and show that the experimental results for electron transport in superfluid $^3\text{He-A}$ provide direct evidence for the spectrum of Weyl Fermions in $^3\text{He-A}$ [5]. I conclude with a discussion of BTRP in chiral superconductors.

[1] T. M. Rice and M. Sigrist, J. Phys. Cond. Mat., **7**, L643 (1995).

[2] T. Mizushima. et al., J.Phys.Soc. Jpn. **85**, 022001 (2016).

[3] H. Ikegami, Y. Tsutsumi, and K. Kono, Science **341**, 59 (2013).

[4] E. R. Schemm, et al. Science **345**, 190 (2014).

[5] O. Shevtsov and J. A. Sauls, Phys. Rev. B **96**, 064511 (2016).